

Tabla 3: Perfil medio de LDL-ácidos grasos (% de ácidos grasos totales) inicial (T0), luego de la dieta de ajuste (T1), luego de la dieta con aceite de maíz (AM), y de la dieta con la mezcla de aceites (MA) (20)

% de ácidos grasos totales	T1 (luego de la dieta de ajuste)	grupo AM (aceite de maíz)	T2 (mezcla de aceites)	grupo MA
C14:0	1.04 ± 0.3	0.85 ± 0.3		1.05 ± 0.3
C16:0	18.95 ± 2.1	16.99 ± 2.4		18.75 ± 2.1
C16:1n7	1.57 ± 0.5	0.88 ± 0.5		1.38 ± 0.4
C18:0	7.28 ± 0.7	6.99 ± 0.8		7.12 ± 0.9
C18:1n9	17.71 ± 1.8	15.01 ± 2.5		22.63 ± 2.9
C18:2n6	38.49 ± 2.1	47.41 ± 3.4		36.43 ± 2.9
C18:3n3	0.95 ± 0.2	0.49 ± 0.1		0.43 ± 0.1
C20:3n6	1.47 ± 0.3	1.56 ± 0.3		1.97 ± 0.3
C20:4n6	7.27 ± 1.2	6.03 ± 1.4		6.45 ± 1.4
C20:5n3	2.48 ± 0.2	2.01 ± 0.3		2.02 ± 0.2
C22:4n6	2.04 ± 0.3	1.32 ± 0.4		1.26 ± 0.4
C22:6n3	0.76 ± 0.3	0.63 ± 0.2		0.67 ± 0.2
Total	100 ± 0.0	100 ± 0.0		100 ± 0.0
Relación media entre ácido oleico/linoleico	0.46	0.32		0.62
AGS/AGMI/AGPI	27/20/53	25/16/59		27/25/48