

Tabla 1. Hallazgos clínicos en 155 casos de enfermedad de Kawasaki registrados en México (1977 – 2010)

EDAD – SEXO SIGNOS CLINICOS	Vizcaíno A	González GN	Rodríguez HR	Frenkel SM	Sotelo CN*	Quezada ChG	Del Ángel A	Coria LJ	Gil VL	Reporte Casos Aislados	TOTAL
EDADES	5/12 - 11 ^a X = 3a	9/12 - 9 ^a X = 3a	15/12 - 9 ^a X = 5a	8/12 - 18 ^a X = 4a	5/12 - 15 ^a X=4.7	15/12 - 11 ^a X = 3a	9/12 - 12 ^a X = 3.0 a	2/12 -9 ^a X= 2.7 a	0-16 ^a X=2.7a	2/12 - 9 ^a X= 5 ^a	3 ^a -5 ^a
SEXO	4.3:1	3.3:1	M 1.5:1	M 1.6:1	M 8.3:1	M 2.4:1	M 2.7:1	M 4:1	1.4:1	M 6:1	M 3.3:1
Fiebre>5 días	16	13	10	8	28	17	15	15	22	11	155 (100%)
Inyección Conjuntival	15	13	10	5	28	15	14	9	18	11	138 (89%)
Exantema no vesicular	16	13	10	8	28	17	15	15	20	11	153 (98%)
Cambios en labios	12	13	10	8	28	17	14	10	12	11	139 (89%)
Enantema	10	13	10	8	28	17	11	8	12	11	128(82%)
Lengua en fresa	10	13	10	8	28	17	10	7	10	11	124 (80%)
Eritema y edema en extremidades	15	13	10	8	26	14	12	10	21	11	140 (90%)
Descamación en piel dedos	13	12	8	8	25	14	8	5	12	9	115 (74%)
Eritema área del pañal	-	-	1	-	3	-	4	4	2	5	19 (12%)
Adenitis cervical	5	13	3	7	18	16	5	4	-	6	77 (50%)
Tos	-	-	-	1	11	-	-	4	10	2	28 (18%)
Dolor abdominal	1	-	7	2	6	-	-	4	4	1	25 (16%)
Náusea –Vómito	1	-	6	-	-	-	-	3	5	1	16 (10%)
Hepatomegalia	2	10	3	1	10	-	-	-	14	-	40 (19%)
Esplenomegalia	-	2	-	4	-	-	-	-	-	-	6 (3.8%)
Irritabilidad	-	1	-	2	2	-	-	-	5	-	10 (6.4%)
Ictericia	-	-	-	1	-	-	-	1	-	1	3 (1.9%)
Meningismo	1	1	-	-	2	-	-	-	-	1	5 (3.2%)
Convulsiones	-	-	-	-	3	-	-	1	-	-	4 (2.5%)
Dolor articular	3	7	6	2	5	-	5	2	5	2	37 (23%)
Soplo precordial	-	-	-	2	3	-	-	-	-	4	9 (5.8%)
Disuria	3	-	1	3	5	-	-	1	-	1	14 (9.0%)
Cicatriz de BCG (activado)	-	-	-	-	1	-	-	1	4	2	8 (5.1%)
Diarrea	1	7	-	-	-	-	-	3	4	2	17 (10.9%)

* Archivos de Cardiología 2007; VI Reunión Infectología Colegio de Pediatras de Sonora, Navojoa Son México, Marzo 2010